OTTAWA COUNTY **Drug Overdose Facts**

DRUG OVERDOSE FAST FACTS*

Ottawa County Data (2016-2020)

- There were 20 unintentional drug overdose deaths in Ottawa County.
- Ottawa County had the 49th highest death rate in the state.
- Adults aged 35-54 had the highest rate of death.
- Nearly two-thirds (65%) of people who died had a history of substance abuse.
- Learn the warning signs of an overdose 50% of deaths occurred at a home or apartment.

*Rates per 100,000 population. Some rates are based on a small number of cases and should be interpreted with caution.

Most common substances in **Ottawa County** overdose deaths:

Methamphetamine

Fentanyl

Alcohol

Methadone

Oxycodone

NUMBER OF UNINTENTIONAL OVERDOSE DEATHS BY TYPE OF SUBSTANCE

OTTAWA COUNTY, 2016-2020

- All unintentional drug overdoses
- Methamphetamine
- **Opioids**
- Benzodiazepines
- Cocaine



12 Oklahomans die every week from an unintentional drug overdose

STATE FACTS:

- From 2016-2020, more than 3,300 Oklahomans died of an unintentional drug overdose.
- From 2007-2020, the unintentional methamphetaminerelated overdose death rate increased 12-fold. In 2020, methamphetamine was involved in nearly two out of three (64%) unintentional drug overdose deaths.
- From 2013-2019, the rate of unintentional prescription opioidrelated overdose death decreased by 58% (501 deaths in 2013; 215 deaths in 2019).
- From 2018-2020, nearly nine out of ten (87%) drug overdose deaths were unintentional and 8% were due to suicide.



Drug Overdose Hospitalizations*

(2017-2019)

Oklahoma State Facts

■ From 2017-2019, more than 14,000 Oklahomans were hospitalized for a nonfatal drug overdose. That's more than 90 Oklahomans each week.

Oklahomans are hospitalized each week for a drug overdose

- Nearly half (49%) of hospitalizations were unintentional and 48% were due to self harm.
- Females were 51% more likely to be hospitalized than males.
- Adults aged 50-59 had the highest rates of hospitalization.
- The most common substances involved in hospitalizations were:
 - Opioids (22%)
 - Benzodiazepines (17%)
 - Non-opioid pain medications (13%)
 - Antidepressants (13%)
 - Stimulants (12%)



Females were 51% more likely to be hospitalized than males.



Ottawa County Facts

- Ottawa County had the 32nd highest nonfatal drug overdose hospitalization rate in the state.
- 45% of hospitalizations were unintentional and 54% due to self harm.
- Females were 32% more likely to be hospitalized than males.
- Adults aged 35-44 had the highest hospitalization rates.
- The most common substances involved in hospitalizations were:
 - Opioids (16%)
 - Antipsychotics (11%)
 - Stimulants (9%)
 - Non-opioid pain medications (9%)
 - Antidepressants (8%)

*Includes all intents of nonfatal drug overdose – unintentional, self harm, undetermined, and assault

Unintentional Drug Overdose Death Rates by County of Residence

Oklahoma, 2016-2020 Grant Rates per 100,000 population Top 5 Counties Dewey Blaine Kinafisher Adair 22.4-28.7 Roger Mills 17.6-22.3 12.3-17.5 Cleveland Caddo Grady 5.8-12.2 Kiowa <5 deaths Coal State Rate: 16.5 Cotton Data Source: Fatal Unintentional Poisoning Surveillance System McCurtain Compiled by: OSDH Injury Prevention Service County of residence was unknown for 43 decedents. Some rates are based on small numbers and may be unstable.

